

JANUARY-2022

A Complete Range of
Diabetic & Cardiac Care

AN
ENDEAVOR
TO PUT LIFE IN LIFE



Perfect
Remedies®

AN ETHICAL DIVISION

AN ASSOCIATE COMPANY OF SOLITAIRE GROUP OF COMPANIES

WHO-GMP CERTIFIED COMPANY

| S.No. | FORMULA <small>{ Read Alphabetically }</small> | BRAND NAME | Where to use INDICATIONS and DOSAGE |
|----------|---|--|--|
| 1. | Alfuzocin HCl. (ER) 10 mg. + Dutasteride 0.5 mg | TABLET DST-AZ | Indications : For Benign prostatic hypertrophy (BPH) in Male with Enlarge Prostate Dosage : 1 tablet daily |
| 2. 3. | Ambrisentan $\frac{5 \text{ mg}}{10 \text{ mg}}$ | TABLET PULMOCOOL $\frac{5}{10}$ | Indications : For the treatment of pulmonary arterial hypertension . Dosage : 5 mg once daily. May increase to 10 mg. if necessary. |
| 4. | Amiloride 5 mg + Hydrochlorothiazide 50 mg | TABLET DILMIL-H | Indications : High blood pressure (hypertension), Heart failure, or extra fluid in the body (edema). Lowering high blood pressure, Helps prevent strokes, Heart attacks, and Kidney problems. Dosage : 1 tablet daily. |
| 5. 6. | Amiloride $\frac{5 \text{ mg}}{20 \text{ mg}}$ & $\frac{5 \text{ mg}}{40 \text{ mg}}$ Frusemide | TABLET DILMIL-FM $\frac{25}{45}$ | Indications : High blood pressure (hypertension), Heart failure, or extra fluid in the body (edema). Lowering high blood pressure, Helps prevent strokes, Heart attacks, and Kidney problems. Dosage : 1 tablet daily. |
| 7. | Amiloride 5 mg + Torsemide 10 mg | TABLET DILMIL-TM | Indication : Edema Associated with congestive heart failure & hypertension Dosage : Once a day |
| 8. 9. | Amlodipine (as besylate) $\frac{5 \text{ mg.}}{10 \text{ mg.}}$ | TABLET WELDIP $\frac{5}{10}$ ✓✓ | Indications : Hypertension, Coronary Artery Disease, Chronic Stable Angina, Vasospastic Angina (Prinzmetal's or Variant Angina), Angiographically Documented Coronary Artery Disease in patients without heart failure or an ejection fraction < 40% Dosage : 5 mg once daily. max. dose 10 mg daily. |
| 10. | Amlodipine (as besylate) 5 mg. + Atenolol 50 mg. | TABLET WELDIP-AT ✓ | Indication : Patients with essential hypertension Dosage : One tablet daily. Max: 2 tablet daily |
| 11. | Amlodipine (as besylate) 5 mg. + Hydrochlorothiazide 12.5 mg. | TABLET WELDIP-H-5 | Indications : Hypertension; Dosage : One tablet once daily. Max: 2 tablet daily. |
| 12. | Amlodipine (as besylate) 10 mg. + Hydrochlorothiazide 12.5 mg. | TABLET WELDIP-H-10 | Indications : Hypertension; Dosage : One tablet once daily. Max: 2 tablet daily. |
| 13. | Amlodipine (as besylate) 10 mg + Indapamide 1.5 mg. (SR) | TABLET WELDIP-IPM | Indications : Hypertension; Dosage : One tablet once daily. Max: 2 tablet daily. |
| 14. | Amlodipine (as besylate) 2.5 mg + Lisinopril 2.5 mg. | TABLET WELDIP-L-2.5 | Indications : Hypertension, Heart Failure, Acute Myocardial Infarction. Dosage : One Tablet Daily. The Dosage may be increased to two tablets daily. The Dosage however should be Individualized. |
| 15. | Amlodipine (as besylate) 5 mg + Lisinopril 5 mg. | TABLET WELDIP-L-5 | Indications : Hypertension, Heart Failure, Acute Myocardial Infarction. Dosage : One Tablet Daily. The Dosage may be increased to two tablets daily. The Dosage however should be Individualized. |
| 16. | Amlodipine (as besylate) 5 mg. + Metoprolol 25 mg. <small>(Metoprolol Succinate equi. to Tartarate)</small> | TABLET WELDIP-AM-30 | Indications : Hypertension, Angina pectoris. Dosage : One tablet once daily. Max: 2 tablet daily. |
| 17. | Amlodipine (as besylate) 5 mg. + Metoprolol 50 mg. <small>(Metoprolol Succinate equi. to Tartarate)</small> | TABLET WELDIP-AM-55 ✓ | Indications : Hypertension, Angina pectoris. Dosage : One tablet once daily. Max: 2 tablet daily. |
| 18. | Amlodipine (As besylate) 5 mg + Ramipril 2.5 mg. | TABLET WELDIP-R-7.5 | Indication : Hypertension Dosage : One tablet daily. Max: 2 tablet daily |

ABBREVIATIONS :

DT = SWEET FLAVOURED DISPERSIBLE TABLET SR / CR / ER = SUSTAINED / CONTROLLED / EXTENDED RELEASE MD = SWEET FLAVOURED MOUTH DISSOLVING
 - AVAILABLE PRODUCTS ARE MARKED WITH TICK (✓) REST PRODUCTS ARE BEING MADE AVAILABLE ON MARKET DEMAND TIME TO TIME
 - BILAYERED = 2 COLOUR TABLET - (MEANS FRONT SIDE ONE COLOUR & BACK SIDE 2ND COLOUR)

| S.No. | FORMULA <small>{ Read Alphabetically }</small> | BRAND NAME | Where to use INDICATIONS and DOSAGE |
|-------------------|---|--|--|
| 19. | Aspirin 75 mg. + Clopidogrel 150 mg. (as bisulphate) | TABLET CLOPITOP-225 ✓ | Indication : For Prevention of ischemic events, myocardial infarction, stroke and cardiovascular death in patients with acute coronary syndrome. Dosage : one tablet daily. |
| 20. | Aspirin 75 mg. + Clopidogrel 75 mg. (as bisulphate) | TABLET CLOPITOP-150 | Indication : For Prevention of ischemic events, myocardial infarction, stroke and cardiovascular death in patients with acute coronary syndrome. Dosage : one tablet daily. |
| 21. | Aspirin 150 mg + Glycine 75 mg | TABLET GLYPRIN-AG | Indications : Conditions where modification of platelet behaviour is considered beneficial, including transient ischaemic attacks, secondary prevention of myocardial infarction, and for prophylaxis against stroke, vascular occlusion and deep vein thrombosis. Dosage : Adults: 1 tab daily or as required. |
| 22. | Atenolol 50 mg. | TABLET 50 -AT ✓ | Indications : Chest pain (angina) & high blood pressure. Dosage : 50 mg per day given once daily. Max: 100mg/day |
| 23. | Atenolol 50 mg. + Chlorthalidone 6.25 mg. | TABLET 50-ATC | Indications : Essential Hypertension Dosage : One - tablet once daily. |
| 24. | Atenolol 50 mg + Losartan Potassium 50 mg. | TABLET 50 -ATL | Indications : Reduce the risk of stroke in patients with hypertension and left ventricular hypertrophy. Dosage : One tablet once daily. Max: 2 tablet once daily |
| 25. 26. 27. | Atorvastatin $\frac{10 \text{ mg.}}{20 \text{ mg.}}$ $\frac{80 \text{ mg.}}$ | TABLET ETOVAS $\frac{10}{20}$ ✓ $\frac{80}{80}$ | {Anti-Hyperlipidemia} Indications : Primary hypercholesterolemia (heterozygous familial and non familial), mixed (combined) dyslipidemia & homozygous hypercholesterolemia in patients who have not responded adequately to diet & other appropriate measures. Dosage : 10 - 80 mg OD |
| 28. | Atorvastatin 10 mg + Amlodipine 5 mg. | TABLET ETOVAS-AM-10 | {For Co-Existing Hypertension & Hyperlipidemia} Indications : Hypertension, Chronic Stable Angina, Vasospastic Angina Dosage : One tablet daily. |
| 29. | Atorvastatin 20 mg + Amlodipine 10 mg. | TABLET ETOVAS-AM-20 | {For Co-Existing Hypertension & Hyperlipidemia} Indications : Hypertension, Chronic Stable Angina, Vasospastic Angina Dosage : One tablet daily. |
| 30. | Atorvastatin 40 mg + Amlodipine 10 mg. | TABLET ETOVAS-AM-40 | {For Co-Existing Hypertension & Hyperlipidemia} Indications : Hypertension, Chronic Stable Angina, Vasospastic Angina Dosage : One tablet daily. |
| 31. | Atorvastatin 10 mg + Aspirin 150 mg. | TABLET ETOVAS-AS | Indications : For the treatment of dyslipidemia associated with atherosclerotic arterial disease with risk of Myocardial infarction, Stroke or Peripheral angiopathy. Dosage : One tablet once or twice daily . |
| 32. | Atorvastatin 10 mg. + Clopidogrel 75 mg. | TABLET ETOVAS-CDG ✓ | Indications : For treatment of patients with both hypertension & hypercholesterolemia. Dosage : 1 tablet daily |
| 33. | Atorvastatin Calcium 10 mg + Ezetimibe 10 mg | TABLET ETOVAS-EZ | Indications : As adjunctive therapy to diet for the reduction of elevated total cholesterol, LDL and Apo B in patients with primary (heterozygous familial & non-familial) hypercholesterolemia. Dosage : One tablet once daily |
| 34. | Atorvastatin Calcium 20 mg + Ezetimibe 10 mg | TABLET ETOVAS-EZ-PLUS | Indications : As adjunctive therapy to diet for the reduction of elevated total cholesterol, LDL and Apo B in patients with primary (heterozygous familial & non-familial) hypercholesterolemia. Dosage : One tablet once daily |

ABBREVIATIONS :

DT = SWEET FLAVOURED DISPERSIBLE TABLET SR / CR / ER = SUSTAINED / CONTROLLED / EXTENDED RELEASE MD = SWEET FLAVOURED MOUTH DISSOLVING
 - AVAILABLE PRODUCTS ARE MARKED WITH TICK (✓) REST PRODUCTS ARE BEING MADE AVAILABLE ON MARKET DEMAND TIME TO TIME
 - BILAYERED = 2 COLOUR TABLET - (MEANS FRONT SIDE ONE COLOUR & BACK SIDE 2ND COLOUR)

| S.No. | FORMULA <small>{ Read Alphabetically }</small> | BRAND NAME | Where to use INDICATIONS and DOSAGE |
|-------------------|--|---|--|
| 35. | Atorvastatin Calcium 40 mg + Ezetimibe 10 mg | TABLET ETOVAS-EZ FORTE | Indications : As adjunctive therapy to diet for the reduction of elevated total cholesterol, LDL and Apo B in patients with primary (heterozygous familial & non-familial) hypercholesterolemia. Dosage : One tablet once daily |
| 36. | Atorvastatin 10 mg + Fenofibrate 160 mg | TABLET ETOVAS-FB ✓ | Indications : As an adjunct to diet to reduce elevated total-C, LDL-C, apo B, and TG levels & to increase HDL-C in these patients. Dosage : one tablet once daily. |
| 37. 38. | Atorvastatin $\frac{10 \text{ mg.}}{25 \text{ mg. (ER)}}$ & $\frac{10 \text{ mg.}}{50 \text{ mg. (ER)}}$ Metoprolol | TABLET ETOVAS-MTL $\frac{25}{50}$ | Indication & Dosage : For hypertension : 1 tablet once daily; may increase to 2 tablets daily if the BP control is inadequate after 1-2 wk. |
| 39. | Atorvastatin Calcium 10 mg + Ramipril 2.5 mg | TABLET ETOVAS-R-12.5 | {For Hypertension & Hypercholesterolemia} Indications : Chronic Angina Pectoris, hypertension, myocardial infarction, myocardial re-infarction prevention. Dosage : One tablet daily. the dosage may be increased to two tablets daily. |
| 40. | Atorvastatin Calcium 10 mg + Ramipril 5 mg | TABLET ETOVAS-R-15 | {For Hypertension & Hypercholesterolemia} Indications : Chronic Angina Pectoris, hypertension, myocardial infarction, myocardial re-infarction prevention. Dosage : One tablet daily. the dosage may be increased to two tablets daily. |
| 41. | Atorvastatin 10 mg. + Ramipril 2.5 mg. + Metoprolol 25 mg. {Metoprolol Succinate (eq. to Metoprolol Tartrate)} | CAPSULE ETOVAS-RM-37.5 | Indications : As 2nd line treatment of patients with both essential hypertension and hypercholesterolemia who are not adequately controlled by combination of an anti-hypertensive with lipid lowering agent. Dosage : 1 Capsule daily dose should be titrated as per the requirement |
| 42. | Atorvastatin 10 mg. + Ramipril 5 mg. + Metoprolol 50 mg. {Metoprolol Succinate (eq. to Metoprolol Tartrate)} | CAPSULE ETOVAS-RM- 65 | Indications : As 2nd line treatment of patients with both essential hypertension and hypercholesterolemia who are not adequately controlled by combination of an anti-hypertensive with lipid lowering agent. Dosage : 1 Capsule daily dose should be titrated as per the requirement |
| 43. | Atorvastatin 20 mg. + Ramipril 10 mg. + Metoprolol 50 mg. {Metoprolol Succinate (eq. to Metoprolol Tartrate)} | CAPSULE ETOVAS-RM- 80 | Indications : As 2nd line treatment of patients with both essential hypertension and hypercholesterolemia who are not adequately controlled by combination of an anti-hypertensive with lipid lowering agent. Dosage : 1 Capsule daily dose should be titrated as per the requirement |
| 44. | 3C { Calcium Aspartate 500 mg. + Calcium Orotate 500 mg. + Calcitriol 0.25 mcg. + M { Methylcobalamin 1500 mcg. + Magnesium Hydroxide 50 mg. + • Levocarnitine 150 mg. + V { Vitamin E 200 IU + Folic Acid 1.5 mg. (Vit. B9) + • Zinc Sulphate 7.5 mg. | TABLET 3CM-LVZ L = Levocarnitine V = Vitamins Z = Zinc | Indication : Osteoporosis, Osteopenia, Improves Muscle Strength, Osteo Arthritis, Fractures Dosage : One tablet daily. Preferably with meal |
| 45. 46. 47. | Carvedilol $\frac{3.125 \text{ mg.}}{6.25 \text{ mg.}}$ $\frac{12.5 \text{ mg.}}$ | TABLET CARVOCARE $\frac{3.125}{6.25}$ $\frac{12.5}{12.5}$ | Indications : Heart failure, Hypertension, Left ventricular dysfunction following myocardial infarction. Dosage : 1 tablet twice daily; dose should be titrated as per the requirement |
| 48. 49. 50. | Carvedilol $\frac{10 \text{ ER}}{20 \text{ ER}}$ $\frac{40 \text{ ER}}$ | TABLET CARVOCARE $\frac{10}{20}$ $\frac{40}{40}$ | Indications : Heart failure, Hypertension, Left ventricular dysfunction following myocardial infarction. Dosage : 1 tablet twice daily; dose should be titrated as per the requirement |
| 51. 52. | Chlorthalidone $\frac{6.25 \text{ mg.}}{12.5 \text{ mg.}}$ | TABLET MYCTD $\frac{6.25}{12.5}$ | Indications : Hypertension, Edema & Heart Failure Dosage : 6.25 mg. - 200 mg. per day. Dose should be titrated as per the requirement |

ABBREVIATIONS :

DT = SWEET FLAVOURED DISPERSIBLE TABLET SR / CR / ER = SUSTAINED / CONTROLLED / EXTENDED RELEASE MD = SWEET FLAVOURED MOUTH DISSOLVING
- AVAILABLE PRODUCTS ARE MARKED WITH TICK (✓) REST PRODUCTS ARE BEING MADE AVAILABLE ON MARKET DEMAND TIME TO TIME
- BILAYERED = 2 COLOUR TABLET - (MEANS FRONT SIDE ONE COLOUR & BACK SIDE 2ND COLOUR)

| S.No. | FORMULA <small>{ Read Alphabetically }</small> | BRAND NAME | Where to use INDICATIONS and DOSAGE |
|-------------------|---|--|--|
| 53. 54 | Cilnidipine 5 mg. & 10 mg. | TABLET CILCIL $\frac{5}{10}$ ✓✓ | Indications : Hypertension Dosage : Adult: 5-10 mg once daily, increase to 20 mg once daily if necessary. |
| 55. 56. 57. | Cinacalcet HCl $\frac{30 \text{ mg.}}{60 \text{ mg.}}$ $\frac{60 \text{ mg.}}{90 \text{ mg.}}$ | TABLET CINAMORE $\frac{30}{60}$ $\frac{60}{90}$ | Indications : For the treatment of secondary hyperparathyroidism in patients with chronic kidney diseases (CKD) on dialysis Dosage : 30 mg once daily to max: 180mg once daily & For hypercalcemia in patients with parathyroid carcinoma : 30 mg twice daily to max: 90 mg 3-4 times daily. |
| 58. 59. | Doxazosin Mesylate $\frac{1 \text{ mg.}}{2 \text{ mg.}}$ | TABLET DOXAZO $\frac{1}{2}$ | Indication & Dosage : Benign Prostatic Hyperplasia : 1 - 8 mg Once Daily Hypertension : 1-16 mg Once Daily. |
| 60. | Doxazosin Mesylate 4 mg. (ER) (Extended Release) | TABLET DOXAZO-4 ER | Indication & Dosage : Benign Prostatic Hyperplasia : 1 - 8 mg Once Daily Hypertension : 1-16 mg Once Daily. |
| 61. | Dronedaron (as Hydrochloride) 400 mg. | TABLET DRONCARD-400 | Indications : To reduce the risk of cardiovascular hospitalization In patients with paroxysmal or persistent atrial fibrillation (AF) or atrial flutter (AFL), with a recent episode of AF/AFL & associated cardiovascular risk factor. Dosage : One tablet twice a day with morning & evening meals |
| 62. 63. | Enoxaperine $\frac{40 \text{ mg. / 0.4 mL.}}{60 \text{ mg. / 0.6 mL.}}$ PRE-FILLED-SYRINGE | INJECTION WOXAPIN $\frac{40}{60}$ | Indication : Deep Vein Thrombosis, Prophylaxis, Abdominal surgery, Knee or Hip Replacement Surgery, Medical patients with restricted mobility . Dosage : 40 mg once daily administered by SC injection with the initial dose given 2 hours prior to surgery. The usual duration of administration is 7 to 10 days; up to 12 days administration has been administered in clinical trials. |
| 64. | Epalrestat 50 mg | TABLET EPALFIT-50 ✓ | Indications : For the Improvement of subjective neuropathy symptoms, abnormality of vibration sense, and abnormal changes in heart beat associated with diabetic peripheral neuropathy Dosage : One tablet (50mg) was orally administered three times daily, before each meal. |
| 65. | Epalrestat 50 mg. + Methylcobalamin 500 mcg. | TABLET EPALFIT-M | Indications : Diabetic neuropathy and other microvascular complication, retinopathy / nephropathy, cardio protection in diabetic patients. Dosage : 50 mg 3 times/day before each meal. (dose calculated on the bases of epalrestat dosing.) |
| 66. 67. | Eprosartan Mesylate $\frac{400 \text{ mg}}{600 \text{ mg}}$ | TABLET EPROVISA $\frac{400}{600}$ | Indication : Hypertension. Dosage : Initial: 600 mg per Day Maintenance: 400-800 mg/day once or in two divided doses |
| 68. | Ezetimibe 10 mg + Fenofibrate 160 mg. | TABLET FIBATIME-EZ | Indications : For the treatment of Combined Hyperlipidemia in patients with normal hepatic and renal function Dosage : One tablet once daily. |
| 69. 70. | Felodipine $\frac{5 \text{ mg}}{10 \text{ mg}}$ (Prolonged Release) | TABLET FELODAY $\frac{5}{10}$ | Indications : Hypertension, Chronic Stable Angina Pectoris Dosage : 5 mg once daily. May increase to 10 mg. if necessary. |
| 71. 72. | Furosemide $\frac{50 \text{ mg.}}{100 \text{ mg.}}$ | TABLET FRUSSY $\frac{50}{100}$ | Indications : For the treatment of pulmonary arterial hypertension, Edema, Cognitive Heart Failure & Hypertension Dosage : 1 Tablet once daily. Max. to 2 Tablets daily |
| 73. 74. | Furosemide $\frac{20 \text{ mg.}}{5 \text{ mg.}}$ & $\frac{40 \text{ mg.}}{5 \text{ mg.}}$ Amiloride | TABLET FRUSSY-AR $\frac{25}{45}$ | Indications : For the treatment of pulmonary arterial hypertension, Edema, Cognitive Heart Failure & Hypertension Dosage : 1 Tablet once daily. Max. to 2 Tablets daily |

ABBREVIATIONS :

DT = SWEET FLAVOURED DISPERSIBLE TABLET SR / CR / ER = SUSTAINED / CONTROLLED / EXTENDED RELEASE MD = SWEET FLAVOURED MOUTH DISSOLVING

- AVAILABLE PRODUCTS ARE MARKED WITH TICK (✓) REST PRODUCTS ARE BEING MADE AVAILABLE ON MARKET DEMAND TIME TO TIME

- BILAYERED = 2 COLOUR TABLET - (MEANS FRONT SIDE ONE COLOUR & BACK SIDE 2ND COLOUR)

| S.No. | FORMULA <small>{ Read Alphabetically }</small> | BRAND NAME | Where to use INDICATIONS and DOSAGE |
|-------------------|--|---|--|
| 75. | Furosemide 20 mg. + Potassium chloride 300 mg. | TABLET FRUSSY-KC | Indications : Patients requiring diuresis and concomitant potassium supplementation as in cardiac, pulmonary, hepatic, renal & peripheral edema of various etiologies. Dosage : 1 tab daily preferably in the morning. |
| 76. | Frusamide 20 mg. + Spironolactone 50 mg. | TABLET FRUSSY-SL-25 | Indications : Management of essential hypertension should be restricted to patients with demonstrated hyperaldosteronism. Dosage : 1 - 4 tablet daily. |
| 77. | Frusamide 40 mg. + Spironolactone 50 mg. | TABLET FRUSSY-SL-50 | Indications : Management of essential hypertension should be restricted to patients with demonstrated hyperaldosteronism. Dosage : 1 - 4 tablet daily. |
| 78. 79. | Heparin Sodium $\frac{25000 \text{ I.U.}}{5000 \text{ I.U.}}$ | INJECTION HEPACARD $\frac{25K}{5K}$ | Indications & Dosage : Thrombophlebitis, pulmonary embolism & occlusive vascular disease. It is also used to prevent thromboembolic complications arising from cardiac & vascular surgery, frostbite, dialysis and other perfusion procedures. Heparin is also used as an anticoagulant in blood transfusions. Adults: Treatment may be given by the following routes: 1. Continuous intravenous infusion: a bolus dose of 5,000 units may be given initially followed by an infusion of 20,000 to 40,000 units in 1 litre of sodium chloride intravenous infusion or glucose intravenous infusion over 24 hours. 2. Intermittent intravenous injection: an initial dose of 10,000 units followed by 5,000 to 10,000 units every 4 to 6 hours may be given. 3. Deep subcutaneous injection: the usual dose is 5,000 units injected intravenously followed by subcutaneous injection of 10,000 units 8 hourly or 15,000 units 12 hourly. A concentrated form of heparin injection should be used (eg. 25,000 units/mL). Children: A suggested dosage is 50 units/kg bodyweight initially by I.V. infusion followed by 100 units/kg bodyweight every 4 hours according to the clotting time. |
| 80. 81. | Irbesartan $\frac{150 \text{ mg.}}{300 \text{ mg.}}$ | TABLET IRBEPURE $\frac{150}{300}$ | Indications & Dosage : Hypertension 150 mg once daily can be increased to 300mg once daily / Nephropathy in Type 2 Diabetic Patients: starting with 75mg target maintenance dose is 300 mg once daily Hypovolemia: Start at 75 mg once Daily |
| 82. 83. | Irbesartan $\frac{150 \text{ mg.}}{12.5 \text{ mg.}}$ & $\frac{300 \text{ mg.}}{12.5 \text{ mg.}}$ + HCTZ = Hydrochlorothiazide | TABLET IRBEPURE-H $\frac{150}{300}$ | Indications : High blood pressure (hypertension); This drug may also be used to treat heart failure and to help protect the kidneys from damage due to diabetes. Dosage : 150 mg/12.5 mg once daily. initially; if needed, may titrate up to 300 mg/12.5 mg once daily |
| 84. 85. | Ivabradine (HCl.) $\frac{5 \text{ mg}}{7.5 \text{ mg}}$ | TABLET IVATOP $\frac{5}{7.5}$ | Indications & Dosage : Chronic stable angina pectoris in coronary artery disease patients with normal sinus rhythm who are unable to tolerate or has a contraindication to beta-blockers; or used with beta-blockers in patients inadequately controlled with an optimal dose of beta-blockers and whose heart rate is >60 beats/min Initial: 5 mg twice daily. Increase if necessary to 7.5 mg twice daily after 3-4 weeks |
| 86. | Labetalol 100 mg. | TABLET LBT-100 | Indications : Hypertension Dosage : Initial: 100 mg bid, may increase gradually according to patients response to 200-400 mg bid. Max: 2.4 g/day in 2-4 divided doses. |
| 87. 88. 89. | Lisinopril $\frac{5 \text{ mg.}}{10 \text{ mg.}}$ $\frac{20 \text{ mg.}}$ | TABLET OXOPRIL $\frac{5}{10}$ ✓ 20 | Indications : Hypertension. Dosage : One tablet daily. |
| 90. | Lisinopril 5 mg. + Hydrochlorothiazide 12.5 mg. | TABLET OXOPRIL-H-5 | Indications : Hypertension. Dosage : 1 - 2 tablet daily. |

ABBREVIATIONS :

DT = SWEET FLAVOURED DISPERSIBLE TABLET SR / CR / ER = SUSTAINED / CONTROLLED / EXTENDED RELEASE MD = SWEET FLAVOURED MOUTH DISSOLVING
 - AVAILABLE PRODUCTS ARE MARKED WITH TICK (✓) REST PRODUCTS ARE BEING MADE AVAILABLE ON MARKET DEMAND TIME TO TIME
 - BILAYERED = 2 COLOUR TABLET - (MEANS FRONT SIDE ONE COLOUR & BACK SIDE 2ND COLOUR)

| S.No. | FORMULA <small>{ Read Alphabetically }</small> | BRAND NAME | Where to use INDICATIONS and DOSAGE |
|----------------------|--|---|--|
| 91. | Lisinopril 10 mg. + Hydrochlorothiazide 12.5 mg. | TABLET OXOPRIL-H-10 | Indications : Hypertension. Dosage : 1 - 2 tablet daily. |
| 92. 93. | Losartan Potassium $\frac{25 \text{ mg.}}{50 \text{ mg.}}$ | TABLET ETOTAN $\frac{25}{50}$ ✓ | Indications : Management of hypertension (alone or in combination with other classes of antihypertensive agents, including diuretics) Dosage : 25–100 mg daily, given in 1 dose or 2 divided doses |
| 94. | Losartan Potassium 50 mg. + Amlodipine 5 mg. | TABLET ETOTAN-AM ✓ | Indications : Mild to moderate hypertension Dosage : One tablet daily. Max: 2 tablets daily |
| 95. | Losartan Potassium 50 mg. + Atenolol 50 mg. | TABLET ETOTAN-AT ✓ | Indications : Mild to moderate hypertension Dosage : One tablet daily. Max: 2 tablets daily |
| 96. 97. | Losartan $\frac{25 \text{ mg.}}{12.5 \text{ mg.}}$ & $\frac{50 \text{ mg.}}{12.5 \text{ mg.}}$ Chlorthalidone | TABLET ETOTAN-CH $\frac{25}{50}$ ✓ | Indications : For the treatment of mild to moderate hypertension in adult patients whose blood pressure is not adequately controlled by monotherapy . Dosage : 1 tablet once or twice daily. Dose should be titrated depending upon condition and response of patient. |
| 98. | Losartan $\frac{50 \text{ mg.}}{12.5 \text{ mg.}}$ HCTZ HCTZ = Hydrochlorothiazide | TABLET ETOTAN-H-50 ✓ | Indications : Hypertension ; high blood pressure together with an enlargement in the left side of their heart (left ventricular hypertrophy). Dosage : One tablet daily. Max: 2 tablet daily |
| 99. | Losartan Potassium 100 mg + Hydrochlorothiazide 25 mg. | TABLET ETOTAN-H-100 | Indications : Hypertension. Dosage: 1-2 tablet per day. |
| 100. | Losartan 50 mg + Ramipril 2.5 mg | TABLET ETOTAN-R-2.5 | Indications : For Treatment of Hypertension uncontrolled by monotherapy Dosage : 1 tablet daily. |
| 101. | Losartan 50 mg + Ramipril 5 mg | TABLET ETOTAN-R-5 | Indications : For Treatment of Hypertension uncontrolled by monotherapy Dosage : 1 tablet daily. |
| 102. 103. | Metoprolol Succinate $\frac{25 \text{ mg.}}{50 \text{ mg.}}$ (eq. to Metoprolol Tartrate) | TABLET TAKE-MTL $\frac{25}{50}$ ✓✓ | Indication : Hypertension, angina pectoris, Tachycardia Dosage : Start with 25mg bid; max: 200 bid |
| 104. | Metoprolol 25 mg + Chlorthalidone 12.5 mg {Metoprolol Succinate (eq. to Metoprolol Tartrate)} | TABLET (BILAYERED) TAKEMTL-CH-25 | Indications : Mild-to-moderate essential hypertension Dosage : one tablet once or twice daily |
| 105. | Metoprolol 50 mg + Chlorthalidone 12.5 mg {Metoprolol Succinate (eq. to Metoprolol Tartrate)} | TABLET (BILAYERED) TAKEMTL-CH-50 | Indications : Mild-to-moderate essential hypertension Dosage : one tablet once or twice daily |
| 106. 107. 108. | Metolazone $\frac{2.5 \text{ mg.}}{5 \text{ mg.}}$ $\frac{5 \text{ mg.}}{10 \text{ mg.}}$ | TABLET MZO $\frac{2.5}{5}$ $\frac{5}{10}$ | Indications & Dosage : Hypertension Initial:1.25 mg/day, adjusted after 3-4 wk. Usual range: 2.5 - 5 mg/day, either alone or with other Anti-hypertensives . Maintenance: 5 mg on alternate days. Oedema 5-10 mg/day, up to 20 mg/day if needed. Max: 80 mg/24 hr. |
| 109. | Metolazone 2.5 mg + Losartan Pot. 50 mg. | TABLET METOBEST-LST | Indications : For hypertension in patients whose blood pressure is not adequately controlled in mono-therapy Dosage : 1 - 2 tablets daily. Dose titration should be done depending upon response of patient & severity of condition. |
| 110. 111. | Metolazone $\frac{2.5 \text{ mg.}}{2.5 \text{ mg.}}$ & $\frac{2.5 \text{ mg.}}{5 \text{ mg.}}$ + Ramipril | TABLET METOBEST-R $\frac{2.5}{5}$ | Indications : Hypertension, Hypertension With Left Ventricular Hypertrophy. Dosage : One Tablet Once Daily |

ABBREVIATIONS :

DT = SWEET FLAVOURED DISPERSIBLE TABLET SR / CR / ER = SUSTAINED / CONTROLLED / EXTENDED RELEASE MD = SWEET FLAVOURED MOUTH DISSOLVING

- AVAILABLE PRODUCTS ARE MARKED WITH TICK (✓) REST PRODUCTS ARE BEING MADE AVAILABLE ON MARKET DEMAND TIME TO TIME

- BILAYERED = 2 COLOUR TABLET - (MEANS FRONT SIDE ONE COLOUR & BACK SIDE 2ND COLOUR)

| S.No. | FORMULA <small>{ Read Alphabetically }</small> | BRAND NAME | Where to use INDICATIONS and DOSAGE |
|----------------------|--|--|--|
| 112. 113. | Metolazone $\frac{2.5\text{mg.}}{50\text{mg.}}$ & $\frac{5\text{mg.}}{50\text{mg.}}$ + Spironolactone | TABLET METOBEST-S $\frac{2.5}{5}$ | Indications : For the treatment of Oedema associated with Congestive Heart failure, Cirrhosis and Nephritic Syndrome Dosage : Once daily, Titrate dose depending upon severity of condition |
| 114. 115. 116. | Moxonidine $\frac{0.2\text{ mg}}{0.3\text{ mg}}$ $\frac{0.3\text{ mg}}{0.4\text{ mg}}$ | TABLET HYPERMOX $\frac{0.2}{0.3}$ $\frac{0.3}{0.4}$ | Indication : For Mild to Moderate Hypertension. Dosage : 0.2mg - 0.4 mg Once Daily |
| 117. 118. | Nebivolol $\frac{5\text{ mg.}}{5\text{ mg.}}$ & $\frac{5\text{ mg.}}{10\text{ mg.}}$ + Amlodipine | TABLET NOBIDAY-AM $\frac{5}{10}$ | Indication : Hypertension. Dosage : one tablet daily. Dose titration should be done depending upon response of patient and severity of condition. |
| 119. 120. 121. | Nicorandil $\frac{5\text{ mg.}}{10\text{ mg.}}$ $\frac{10\text{ mg.}}{20\text{ mg.}}$ (PR) (PR) - Prolonged Release | TABLET TAKERUN $\frac{5}{10}$ ✓ $\frac{10}{20}$ ✓ | Indication : Angina pectoris. Dosage : Adult: 10 mg bid, increase if necessary. Usual dose: 10-20 mg bid. May use 5 mg bid for patients prone to headache. Max dose: 30 mg bid. |
| 122. 123. | Nitroglycerin $\frac{2.6\text{ mg. (CR)}}{6.4\text{ mg. (CR)}}$ (CR) - Controlled Release | TABLET LEONITE $\frac{2.6}{6.4}$ ✓ | Indications : Angina pectoris, acute (treatment); Angina pectoris, acute (prophylaxis); Hypertension; Myocardial infarction Dosage : 2.6mg to 15.6 mg every twice daily |
| 124. 125. 126. | Olmesartan Medoxomil $\frac{10\text{ mg.}}{20\text{ mg.}}$ $\frac{20\text{ mg.}}{40\text{ mg.}}$ | TABLET TAKEFAST $\frac{10}{20}$ ✓ $\frac{20}{40}$ ✓ | Indications : Hypertension, cardiovascular risk reduction Dosage : 10 - 40 mg one tablet daily. |
| 127. 128. | Olmesartan $\frac{20\text{ mg.}}{5\text{ mg.}}$ & $\frac{40\text{ mg.}}{5\text{ mg.}}$ Amlodipine | TABLET TAKEFAST-A $\frac{25}{45}$ ✓ | Indication : Hypertension Dosage : Once or Twice a day. |
| 129. 130. | Olmesartan $\frac{20\text{ mg.}}{10\text{ mg.}}$ & $\frac{40\text{ mg.}}{20\text{ mg.}}$ Atorvastatin | TABLET TAKEFAST-ATS $\frac{30}{60}$ | Indications : For the treatment of hypertension, chronic stable angina and vasospastic angina associated with dislipidemia Dosage : 1 tablet daily |
| 131. 132. | Olmesartan $\frac{20\text{ mg.}}{12.5\text{ mg.}}$ & $\frac{40\text{ mg.}}{25\text{ mg.}}$ + HCTZ (HCTZ = Hydrochlorothiazide) | TABLET TAKEFAST-H $\frac{20}{40}$ ✓ | Indications : Hypertension, Cardiovascular risk reduction Dosage : Once a day. |
| 133. | Olmesartan medoximil $\frac{20\text{ mg.}}{5\text{ mg.}}$ + Ramipril | TABLET TAKEFAST-R-25 | Indication : Hypertension Dosage : Once a day. |
| 134. | Olmesartan medoximil $\frac{40\text{ mg.}}{5\text{ mg.}}$ + Ramipril | TABLET TAKEFAST-R-45 | Indication : Hypertention Dosage : Once a day. |
| 135. | Olmesartan medoximil $\frac{20\text{ mg.}}{25\text{ mg.}}$ + Metoprolol (ER) {Metoprolol Succinate (eq. to Metoprolol Tartrate)} | TABLET TAKEFAST-M-45 | Indication : Hypertention Dosage : Once a day. |
| 136. | Olmesartan 5 mg. + Amlodipine 2.5 mg. + HCTZ 12.5 mg. | TABLET TAKEFAST-AH-7.5 | Indications : Hypertention Dosage : Once or twice a day |
| 137. | Olmesartan 20 mg. + Amlodipine 5 mg. + HCTZ 12.5 mg. | TABLET TAKEFAST-AH-25 | Indications : Hypertention Dosage : Once or twice a day |
| 138. | Olmesartan 40 mg. + Amlodipine 5 mg. + HCTZ 12.5 mg. | TABLET TAKEFAST-AH-45 | Indications : Hypertention Dosage : Once or twice a day |

ABBREVIATIONS :

DT = SWEET FLAVOURED DISPERSIBLE TABLET SR / CR / ER = SUSTAINED / CONTROLLED / EXTENDED RELEASE MD = SWEET FLAVOURED MOUTH DISSOLVING
- AVAILABLE PRODUCTS ARE MARKED WITH TICK (✓) REST PRODUCTS ARE BEING MADE AVAILABLE ON MARKET DEMAND TIME TO TIME
- BILAYERED = 2 COLOUR TABLET - (MEANS FRONT SIDE ONE COLOUR & BACK SIDE 2ND COLOUR)

| S.No. | FORMULA <small>{ Read Alphabetically }</small> | BRAND NAME | Where to use INDICATIONS and DOSAGE |
|----------------------|---|---|--|
| 139. | Olmesartan medoximil 20 mg + Metoprolol (ER) 50 mg {Metoprolol Succinate (eq. to Metoprolol Tartrate)} | TABLET TAKEFAST-M-50 | Indication : Hypertention Dosage : Once a day. |
| 140. | Pitavastatin 1 mg. | TABLET ETOVAS-PTS-1 | Indications : Hypercholesterolemia Dosage : 1 - 4 mg. daily |
| 141. | Pitavastatin 2 mg. | TABLET ETOVAS-PTS-2 | Indications : Hypercholesterolemia Dosage : 1 - 4 mg. daily |
| 142. | Pitavastatin 4 mg. | TABLET ETOVAS-PTS-4 | Indications : Hypercholesterolemia Dosage : 1 - 4 mg. daily |
| 143. 144. | Prasugrel (as HCl.) $\frac{5 \text{ mg.}}{10 \text{ mg.}}$ | TABLET WILPRASU $\frac{5}{10}$ ✓ | Indications : Acute Coronary Syndrome Dosage : 60 mg. loading dose, then 5 - 10 mg. daily |
| 145. 146. 147. | Prazosin $\frac{1 \text{ mg.}}{2 \text{ mg.}} \frac{5 \text{ mg.}}$ | TABLET PRAZOTAB $\frac{1}{2}$ $\frac{5}{5}$ | Indications & Dosage : Hypertension Initial: 0.5 mg bid or tid for 3-7 days, increase to 1 mg bid or tid for the next 3-7 days if tolerated. Max: 20 mg/day in divided doses. Heart failure Initial: 0.5 mg 2-4 times/day. Maintenance: 4-20 mg/day in divided doses. Benign prostatic hyperplasia; Raynaud's syndrome Initial: 0.5 mg bid. Maintenance: ≤2 mg bid. |
| 148. 149. 150. | Ramipril $\frac{2.5 \text{ mg.}}{5 \text{ mg.}} \frac{10 \text{ mg.}}$ | TABLET OXORAM $\frac{2.5}{5}$ $\frac{10}{10}$ ✓ ✓ | Indication : Hypertension, heart failure & left ventricular dysfunction. Dosage : 1.25 mg - 10 mg once daily |
| 151. | Ramipril 2.5 mg + Hydrochlorothiazide 12.5 mg | TABLET OXORAM-H-2.5 ✓ | Indication : Hypertension Dosage : once or twice a daily |
| 152. | Ramipril 5 mg + Hydrochlorothiazide 12.5 mg | TABLET OXORAM-H-5 | Indication : Hypertension Dosage : once or twice a daily |
| 153. | Ramipril 10 mg + Hydrochlorothiazide 12.5 mg | TABLET OXORAM-H-10 | Indication : Hypertension Dosage : once or twice a daily |
| 154. 155. | Ramipril $\frac{2.5 \text{ mg.}}{1.5 \text{ mg.}}$ & $\frac{5 \text{ mg.}}{1.5 \text{ mg.}}$ Indapamide | TABLET OXORAM-IPM $\frac{2.5}{5}$ ✓ | Indication : Patients with essential hypertension Dosage : One tablet daily. Max: 2 tablet daily |
| 156. 157. | Ramipril $\frac{2.5 \text{ mg.}}{25 \text{ mg.}}$ & $\frac{5 \text{ mg.}}{50 \text{ mg.}}$ Metoprolol (ER) {Metoprolol Succinate (eq. to Metoprolol Tartrate)} | TABLET OXORAM-MTL $\frac{27.5}{55}$ | Indication : Hypertension. Dosage : 1 tablet daily, may be increase to max 3 tablet daily. |
| 158. | Strip of Combi-Pack Contains : A) Ramipril 5 mg. + Amlodipine 5 mg. B) Atorvastatin 10 mg. | TABLET OXORAM-A-10 | Indications : Hypertention Dosage : One tablet of each tablet once daily |
| 159. | Strip of Combi-Pack Contains : A) Ramipril 5 mg. + Amlodipine 5 mg. B) Atorvastatin 20 mg. | TABLET OXORAM-A-20 | Indications : Hypertention Dosage : One tablet of each tablet once daily |
| 160. 161. 162. | Ranolazine $\frac{500 \text{ mg. (ER)}}{750 \text{ mg. (ER)}} \frac{1000 \text{ mg. (ER)}}$ | TABLET RANOZEAL $\frac{500}{750}$ $\frac{1000}{1000}$ ✓ ✓ | Indication : Chronic Angina. Dosage: 500 mg BID initially; may increase to 1000 mg BID; do not exceed 2000 mg/day. Dosage : 1-2 tablets daily. |

ABBREVIATIONS :

DT = SWEET FLAVOURED DISPERSIBLE TABLET SR / CR / ER = SUSTAINED / CONTROLLED / EXTENDED RELEASE MD = SWEET FLAVOURED MOUTH DISSOLVING

- AVAILABLE PRODUCTS ARE MARKED WITH TICK (✓) REST PRODUCTS ARE BEING MADE AVAILABLE ON MARKET DEMAND TIME TO TIME

- BILAYERED = 2 COLOUR TABLET - (MEANS FRONT SIDE ONE COLOUR & BACK SIDE 2ND COLOUR)

| S.No. | FORMULA <small>{ Read Alphabetically }</small> | BRAND NAME | Where to use INDICATIONS and DOSAGE |
|------------------------------|---|---|--|
| 163. 164. 165. 166. | Rosuvastatin $\frac{5 \text{ mg.}}{10 \text{ mg.}}$ $\frac{20 \text{ mg.}}{40 \text{ mg.}}$ | TABLET ROFAST $\frac{5}{10}$ $\frac{20}{40}$ ✓ | Indications : Hyperlipidemia and mixed dyslipidemia, primary prevention of cardiovascular diseases Dosage : 5-40mg once daily |
| 167. 168. | Rosuvastatin $\frac{5 \text{ mg.}}{10 \text{ mg.}}$ & $\frac{10 \text{ mg.}}{10 \text{ mg.}}$ Ezetimibe | TABLET ROFAST-EZ $\frac{5}{10}$ | Indications : Hypercholesterolemia Dosage : 1 tablet once daily. |
| 169. | Rosuvastatin $\frac{5 \text{ mg.}}{160 \text{ mg.}}$ + Fenofibrate | TABLET ROFAST-FBT-5 | Indications : As an adjunct to diet to reduce elevated total - Cholesterol, LDL-C (Total Density Low Cholesterol), Apolipoprotein B (ApoB) and triglyceride level. Dosage : 5 mg once daily. May increase to 10 mg. if necessary. |
| 170. 171. | Rosuvastatin $\frac{10 \text{ mg.}}{160 \text{ mg.}}$ & $\frac{20 \text{ mg.}}{160 \text{ mg.}}$ Fenofibrate | TABLET ROFAST-FBT $\frac{10}{20}$ ✓✓ | Indications : As an adjunct to diet to reduce elevated total - Cholesterol, LDL-C (Total Density Low Cholesterol), Apolipoprotein B (ApoB) and triglyceride level. Dosage : 5 mg once daily. May increase to 10 mg. if necessary. |
| 172. 173. 174. | Sildenafil 2 mg., 4 mg. & 8 mg. | CAPSULE SIDOSIN $\frac{2}{4}$ 8 | Indications : Symptomatic treatment of benign prostatic hyperplasia. Dosage : 1 capsule daily. max. dose 8 mg |
| 175. 176. | Simvastatin $\frac{10 \text{ mg.}}{10 \text{ mg.}}$ & $\frac{20 \text{ mg.}}{10 \text{ mg.}}$ + Ezetimibe | TABLET SIMDUE-EZ $\frac{20}{30}$ | Indications : For primary hypercholesterolemia. Dosage : Once or twice a day |
| 177. | Simvastatin 20 mg. + Hydrochlorothiazide 12.5 mg + Ramipril 5 mg. | TABLET SIMADUE-HR | Indications : For treatment of patients with both hypertension & hypercholesterolemia. Dosage : 1 tablet daily. Individualize the dosage. Titration may be needed in some patients |
| 178. 179. 180. 181. | Telmisartan $\frac{10 \text{ mg.}}{20 \text{ mg.}}$ $\frac{40 \text{ mg.}}{80 \text{ mg.}}$ | TABLET TAKETEL $\frac{10}{40}$ $\frac{20}{80}$ ✓✓ | Indication : Hypertension Dosage : Adult: Initial: 10 mg once daily. Max: 80 mg/day |
| 182. 183. | Telmisartan $\frac{40 \text{ mg.}}{5 \text{ mg.}}$ & $\frac{80 \text{ mg.}}{5 \text{ mg.}}$ + Amlodipine | TABLET TAKETEL-AM $\frac{45}{85}$ ✓ | Indication : Hypertension Dosage : One tablet once daily. if necessary, the dose may be increased to two tablets once daily. |
| 184. | Telmisartan 20 mg. + Amlodipine 2.5 mg. + Hydrochlorothiazide 12.5 mg. | TABLET TAKETEL-AHT-20 | } Indications : Hypertension (where monotherapy is not sufficient) Dosage : One tablet once daily. May increase to 2 tablets |
| 185. | Telmisartan 40 mg. + Amlodipine 5 mg. + Hydrochlorothiazide 12.5 mg. | TABLET TAKETEL-AHT-40 | |
| 186. | Telmisartan 80 mg. + Amlodipine 10 mg. + Hydrochlorothiazide 25 mg. | TABLET TAKETEL-AHT-80 ✓ | |
| 187. 188. | Telmisartan $\frac{40 \text{ mg.}}{10 \text{ mg.}}$ & $\frac{20 \text{ mg.}}{10 \text{ mg.}}$ + Atorvastatin (as calcium) | TABLET TAKETEL-ATS $\frac{20}{40}$ | Indication : For Coexisting Hypertension & Hyperlipidemia. Dosage : One -two tablet daily |
| 189. 190. | Telmisartan + $\frac{20 \text{ mg.}}{25 \text{ mg.}}$ & $\frac{40 \text{ mg.}}{50 \text{ mg.}}$ Metoprolol (ER) (ER = Extended Release) | TABLET TAKETEL-MTL $\frac{45}{90}$ ✓✓ | Indication : Hypertension Dosage : One tablet once daily. if necessary, the dose may be increased to two tablets once daily. |
| 191. 192. | Telmisartan $\frac{40 \text{ mg.}}{2.5 \text{ mg.}}$ & $\frac{40 \text{ mg.}}{5 \text{ mg.}}$ Ramipril | TABLET TAKETEL-R $\frac{42.5}{45}$ ✓ | Indication : Hypertension Dosage : One tablet once or twice a day. |
| 193. 194. | Trimetazidine 20 mg & 35 mg. | TABLET TRIMMY $\frac{20}{35}$ | Indications : Angina pectoris & Meniere's disease Dosage : Adult: As conventional tab: 40-60 mg daily |

ABBREVIATIONS :

DT = SWEET FLAVOURED DISPERSIBLE TABLET SR / CR / ER = SUSTAINED / CONTROLLED / EXTENDED RELEASE MD = SWEET FLAVOURED MOUTH DISSOLVING

- AVAILABLE PRODUCTS ARE MARKED WITH TICK (✓) REST PRODUCTS ARE BEING MADE AVAILABLE ON MARKET DEMAND TIME TO TIME

- BILAYERED = 2 COLOUR TABLET - (MEANS FRONT SIDE ONE COLOUR & BACK SIDE 2ND COLOUR)

| S.No. | FORMULA <small>{ Read Alphabetically }</small> | BRAND NAME | Where to use INDICATIONS and DOSAGE |
|----------------------|---|--|--|
| 195. 196. | Acarbose $\frac{25 \text{ mg.}}{500 \text{ mg.}}$ & Metformin $\frac{50 \text{ mg.}}{500 \text{ mg.}}$ + | TABLET DIADOSE-M $\frac{25}{50}$ ✓ | Indications : For the treatment of patients with Type II diabetes mellitus, when diet, Exercise and single agent do not result in adequate glycemic control Dosage : 1 tablets twice daily |
| 197. | Alendronate Sodium (eq. to Alendronic Acid 10 mg.) (Pack of 30 Tablets) | TABLET OSTOSIS-10 ✓ | Diabetics are prone to osteoporosis (Ref. Source : J Bone Miner Res. 2007 Sep;22(9):1317-28.) Prophylaxis of osteoporosis in women: 5–10 mg daily or 35–70 mg weekly. Therapy of osteoporosis in women & men : 10 mg daily or 70 mg weekly. Osteoporosis under corticosteroids: 5–10 mg daily or 35–70 mg weekly in men and premenopausal women or those receiving concomitant HRT. In postmenopausal women not receiving HRT, the recommended dose is 10 mg daily or 70 mg weekly. |
| 198. | Alendronate Sodium (eq. to Alendronate Acid 70 mg.) + Cholecalciferol 70 mcg. (eq. to 2800 IU Vit. D3) (Pack of 4 Tablets) | TABLET OSTOSIS-PLUS ✓ | Indication : For the treatment of osteoporosis in post-menopausal women. Dosage: 70 mg alendronic acid once weekly, taken by mouth with a glass of plain water, at least 30 minutes before any food, beverages or other medication is taken. Remaining days take calcium carbonate 1 tablet daily. |
| 199. | Bromocriptine 0.8 mg. (as mesylate) | TABLET GLYCOBROM - 0.8 | Indication : As an adjunct to diet & exercise to improve glycemic control in adults with type-2 diabetes mellitus. Dosage : Initiated at one tablet (0.8 mg) and increased by one tablet per week until a maximum daily dose of 6 tablets (4.8 mg) |
| 200. | Calcium Dobisilate 500 mg. | TABLET DOBITIME ✓ | Indications : Diabetic Retinopathy and Glaucoma, Diabetic Microangiopathy, Chronic Venous Insufficiency, Hemorrhoids, Myocardial Infarction Dosage : 500 mg, orally 3 times daily. Max: 1000mg bid |
| 201. 202. 203. | Carbimazole $\frac{5 \text{ mg.}}{10 \text{ mg.}}$ $\frac{10 \text{ mg.}}{20 \text{ mg.}}$ | TABLET THYROCLEAN-CM $\frac{5}{10}$ $\frac{10}{20}$ | Indications : Hyperthyroidism, Preparation for thyroidectomy, Adjunct to radio-iodine therapy, Grave's disease. Dosage : Adult: Initially, 20-60 mg daily as a single or divided dose, reduced gradually once euthyroidism is achieved. Maintenance: 5-15 mg daily for at least 1 year or 18 month. Child : 4-12 yr: Initially, 15 mg daily. |
| 204. 205. | Darifenacin $\frac{7.5 \text{ mg. (ER)}}{15 \text{ mg. (ER)}}$ (as hydrobromide) | TABLET DARVISA $\frac{7.5}{15}$ | Indications : For the treatment of overactive bladder with symptoms of urinary urge incontinence, urgency and frequency Dosage : Starting with 7.5mg which may be increased to 15 mg daily. |
| 206. | Dutasteride 0.5 mg | TABLET DST-0.5 ✓ | Indication : Benign prostatic Hyperplasia (BPH) Dosage : One tablet daily |
| 207. | Epalrestat 50 mg. + Methylcobalamin 500 mcg. | TABLET EPALFIT-M | Indications : Diabetic neuropathy and other microvascular complication, retinopathy / nephropathy, cardio protection in diabetic patients. Dosage : 50 mg 3 times/day before each meal. (Dose calculated on the bases of epalrestat dosing.) |
| 208. 209. | Flavoxate HCl. $\frac{100 \text{ mg.}}{200 \text{ mg.}}$ | TABLET GETFLAVO $\frac{100}{200}$ ✓✓ | Indications : Dysuria, Urinary Frequency, Overactive bladder/ urinary incontinence, Nocturia, Suprapubic pain, Urinary Tract Irritation, Urinary urgency. Dosage : 100-200 mg every 6-8hr. |
| 210. | Glutathione 600 mg. | INJECTION GLULIV-6 | Indications : Treatment of alcoholic liver diseases like alcoholic fatty liver, alcoholic liver fibrosis, alcoholic liver cirrhosis & hepatitis. Dosage : In mild to moderate disease- 600 mg to 2.4 g twice daily by intramuscular or slow intravenous injection. |
| 211. | An Ayurvedic Proprietary Medicine For Weight Lose | CAPSULE EVASLIM | Indication : Ayurvedic Dosage : One Capsule twice or thrice a day |
| 212. | Gliclazide 80 mg. | TABLET UROGLID-80 ✓ | Indications : Non-insulin dependent diabetes mellitus Dosage : Commenced with a total daily dose of 40-80 mg; may be increased to a maximum of 320 mg/day depending on the blood glucose response. |

ABBREVIATIONS :

DT = SWEET FLAVOURED DISPERSIBLE TABLET SR / CR / ER = SUSTAINED / CONTROLLED / EXTENDED RELEASE MD = SWEET FLAVOURED MOUTH DISSOLVING

- AVAILABLE PRODUCTS ARE MARKED WITH TICK (✓) REST PRODUCTS ARE BEING MADE AVAILABLE ON MARKET DEMAND TIME TO TIME

- BILAYERED = 2 COLOUR TABLET - (MEANS FRONT SIDE ONE COLOUR & BACK SIDE 2ND COLOUR)

| S.No. | FORMULA <small>{ Read Alphabetically }</small> | BRAND NAME | Where to use INDICATIONS and DOSAGE |
|------------------------------|--|---|---|
| 213. | Gliclazide 80 mg. + Chromium Picolinate 200 mg. | TABLET UROGLID-C | Indications : Non-insulin dependent diabetes mellitus. Dosage : Commenced with a total daily dose of 40-80 mg; may be increased to a maximum of 320 mg/day depending on the blood glucose response. Chromium Picolinate : Optimal daily Allowance (ODA) is 200-600mcg daily. |
| 214. 215. 216. 217. | Glimepiride $\frac{1 \text{ mg.}}{2 \text{ mg.}}$ $\frac{3 \text{ mg.}}{4 \text{ mg.}}$ | TABLET EUROGLIP $\frac{1}{2} \frac{3}{4}$ ✓✓ | Indications : Type 2 Diabetes. Dosage : Initial Therapy is 1-2mg once daily and may be increased to 4 mg daily in maintenance phase. |
| 218. 219. | Glimepiride $\frac{1 \text{ mg.}}{500 \text{ mg.}}$ & $\frac{2 \text{ mg.}}{500 \text{ mg.}}$ + Metformin | TABLET EUROGLIP $\frac{M1}{M2}$ ✓✓ | Indications : Type 2 Diabetes. Dosage : One Tablet Daily. Dosage may be increased to two tablets daily. the dosage however should be individualized. |
| 220. 221. | Glimepiride $\frac{1 \text{ mg.}}{1000 \text{ mg}}$ & $\frac{2 \text{ mg.}}{1000 \text{ mg}}$ + Metformin SR | TABLET EUROGLIP-PLUS $\frac{M1}{M2}$ ✓ BILAYER TABLETS | Indication : Type 2 diabetes mellitus. Dosage : 1-2 tablet daily |
| 222. 223. | Glimepiride 1 & 2 mg + Metformin 500 (SR) + Pioglitazone 15 mg. | TABLET EUROGLIP $\frac{MP1}{MP2}$ ✓✓ BILAYER TABLETS | Indications : Management of type 2 diabetes (NIDDM) Dosage : One - Two tablet daily. Max. 4 Tablets (NIDDM = Non-Insulin-Dependent Diabetes Mellitus) |
| 224. | Glimepiride 1 mg + Pioglitazone 15 mg. | TABLET EUROGLIP-P | Indications : Management of type 2 diabetes (NIDDM) Dosage : One - Two tablet daily. Max. 4 Tablets (NIDDM = Non-Insulin-Dependent Diabetes Mellitus) |
| 225. | Glimepiride 2 mg & Pioglitazone 30 mg. | TABLET EUROGLIP-FORTE ✓ | Indications : Management of type 2 diabetes (NIDDM) Dosage : One tablet daily. (NIDDM = Non-Insulin-Dependent Diabetes Mellitus) |
| 226. | Ibandronic Acid 150 mg. | TABLET OSTOSIS-IB-150 | Indications : For treatment and prevention of osteoporosis in postmenopausal women; increases BMD and reduces vertebral fracture incidence Dosage : 150 mg every month |
| 227. 228. 229. | Levothyroxine $\frac{25 \text{ mcg.}}{50 \text{ mcg.}}$ $\frac{100 \text{ mcg.}}$ | TABLET THYROCLEAN $\frac{25}{50}$ 100 ✓ | Indications : Hypothyroidism, Pituitary TSH Suppression Dosage : single daily dose, preferably one-half to one-hour before breakfast. (dose titration is required as per response of patient.) |
| 230. 231. | Methimazole $\frac{5 \text{ mg.}}{10 \text{ mg.}}$ | TABLET THYROCLEAN $\frac{M5}{M10}$ | Indications : For Hyperthyroidism Dosage : 15 - 60mg in 3 divided dosage |
| 232. 233. 234. | Miglitol $\frac{25 \text{ mg}}{50 \text{ mg}}$ $\frac{100 \text{ mg}}$ | TABLET MIGLIPUR $\frac{25}{50}$ 100 ✓ | Indication : Type 2 diabetes mellitus. Dosage : 1 Tablet every 8hr at meals (with first bite) |
| 235. | Miglitol 25 mg + Metformin HCl 500 mg. | TABLET MIGLIPUR-M 25 ✓ | Indication : Type - 2 Diabetes. Dosage : One tablet twice /thrice daily |
| 236. | Miglitol 50 mg + Metformin HCl 500 mg. (SR) | TABLET MIGLIPUR-M 50 ✓ | Indication : Type - 2 Diabetes. Dosage : One tablet twice /thrice daily |
| 237. | 1) Mecobalamin 1500mcg + 2) Alpha lipoic Acid 100 mg + 3) Vit B6 3mg + 4) Folic Acid 1.5 mg 5) Zinc 15mg + 6) Chromium 60 mcg 7) Selenium 65 mcg + 8) Biotin 100 mcg + 9) Inositol 2 mg | TABLET MECOPURE-9 ✓ | Indications : Diabetic Neuropathies, Peripheral Neuropathies, Vit.B12 deficiency ,Neurological pain, Sensory disturbances, Hyper-Homocysteinemia Dosage : 1 tablet daily |

ABBREVIATIONS :

DT = SWEET FLAVOURED DISPERSIBLE TABLET SR / CR / ER = SUSTAINED / CONTROLLED / EXTENDED RELEASE MD = SWEET FLAVOURED MOUTH DISSOLVING
 - AVAILABLE PRODUCTS ARE MARKED WITH TICK (✓) REST PRODUCTS ARE BEING MADE AVAILABLE ON MARKET DEMAND TIME TO TIME
 - BILAYERED = 2 COLOUR TABLET - (MEANS FRONT SIDE ONE COLOUR & BACK SIDE 2ND COLOUR)

| S.No. | FORMULA <small>{ Read Alphabetically }</small> | BRAND NAME | Where to use INDICATIONS and DOSAGE |
|-------|--|---|---|
| 238. | Mecobalamin 1500mcg + Alpha lipoic acid 100mg + Inositol 100mg + Vit. A 5000IU + Vit B1 10mg + Pyridoxine 1.5mg + Vit E 25IU + Calcium Pantothenate 10mg + Folic Acid 1500mcg + Chromium 200mcg + Selenium 163.6mcg | TABLET MECOPURE-VITAL | Indications : For the treatment of vitamin deficiency states in adults Dosage : 1 tablet daily |
| 239. | Mecobalamin 750 mcg + Benfotiamine 150 mg + Pyridoxine 1.5 mg + Nicotinamide 45 mg | TABLET MECOPURE-BPN | Indications : Dementia,depression, Insomnia, Sciatica Syndrome, Diabetic Neuropathy, Peripheral Neuropathy, Drug Induced Neuropathy Dosage : 1 - 2 tablet daily |
| 240. | Mecobalamin 1000 mcg. + Thiamine HCl. 100 mg. + Pyridoxine HCl. 100 mg. + Nicotinamide 100 mg. + & D-Panthenol 50 mg. | INJECTION / 2 ML. AMPOULE VITACOB-PLUS (WITH DISPOSABLE SYRINGE) | Indications : Peripheral Neuropathies (like Diabetic & Alcoholic Neuropathy, Drug induced neuropathy, Lumbago, Entrapment Neuropathy, Intercostal Neuralgia & other Neuropathies). Dosage : The usual adult dose is one injection, administered intramuscularly or intravenously three times a week. The dose should be adjusted according to the age of the patient and severity of symptoms. |
| 241. | Metadoxine 500 mg | TABLET METAKARE | Indications & Dosage : Acute Alcohol Intoxication - 1 - 2 tablet per day Alcoholic Fatty Liver - 2 tablet per day |
| 242. | Methylcobalamin 500 mcg + Alpha Lipoic Acid 100 mg. + Benfotiamine 100 mg. + Pyridoxine 50 mg. + | TABLET MECOPURE-ABP | Indications : Peripheral Neuropathies (like Diabetic & Alcoholic Neuropathy, Drug induced neuropathy, Lumbago, Entrapment Neuropathy, Intercostal Neuralgia & other Neuropathies). Dosage : One Tablet Once daily |
| 243. | Methylcobalamin 1500 mcg + Benfotiamine 50 mg. + Biotin 5 mg. + Alpha lipoic Acid 200 mg + Pyridoxine 3 mg + Folic acid 1.5 mg | TABLET MECOPURE-BBA | Indications : Peripheral Neuropathies (like Diabetic & Alcoholic Neuropathy, Drug induced neuropathy, Lumbago, Entrapment Neuropathy, Intercostal Neuralgia & other Neuropathies). Dosage : One Tablet Once daily |
| 244. | Methylcobalamin 1500 mcg + Gabapentin (SR) 600 mg | TABLET MECOPURE-GB-600 | Indications : nerve pain conditions, including peripheral neuropathy, trigeminal neuralgia and diabetic neuropathy Dosage : 1 tablet daily. |
| 245. | Methylcobalamin 1500 mcg + Pregabalin (ER) 150 mg | TABLET MECOPURE-PG-150 | Indication : For the treatment of adult patients with peripheral neuropathy. Dosage : 1 Tablet daily. |
| 246. | Metformin 500 mg. | TABLET EXEED-500 ✓ | Indications : Indicated as an adjunct to diet and exercise to improve glycemic control in patients with type 2 diabetes Dosage : 500 mg - 2000 mg. in divided doses |
| 247. | Metformin 1000 mg. (SR) | TABLET EXEED-SR ✓ | Indications : Type 2 diabetes Dosage : One tablet daily. Max: metformin dose is 2000 mg in adults. |
| 248. | Metformin 500 mg. + Alpha Lipoic Acid 200 mg. | TABLET EXEED-A2 | Indications : For the treatment of patients with Type II diabetes mellitus, when diet, Exercise and single agent do not result in adequate glycemic control Dosage : 1 tablets twice daily |
| 249. | Metformin 500 mg (SR) + Atorvastatin 10 mg | TABLET EXEED-ATS | Indications : As adjunctive therapy to diet & exercise in patients with diabetic dyslipidemia Dosage : One Tablet once daily |
| 250. | Metformin HCl (ER) 500mg + Fenofibrate 160 mg | TABLET EXEED-FBT | Indication : For the treatment of Type-II diabetes associated with mixed dyslipidemia. Dosage: 1 Tablet daily. |

ABBREVIATIONS :

DT = SWEET FLAVOURED DISPERSIBLE TABLET SR / CR / ER = SUSTAINED / CONTROLLED / EXTENDED RELEASE MD = SWEET FLAVOURED MOUTH DISSOLVING
- AVAILABLE PRODUCTS ARE MARKED WITH TICK (✓) REST PRODUCTS ARE BEING MADE AVAILABLE ON MARKET DEMAND TIME TO TIME
- BILAYERED = 2 COLOUR TABLET - (MEANS FRONT SIDE ONE COLOUR & BACK SIDE 2ND COLOUR)

| S.No. | FORMULA <small>{ Read Alphabetically }</small> | BRAND NAME | Where to use INDICATIONS and DOSAGE |
|-------|--|---|--|
| 251. | Metformin 500 mg + Glibenclamide 5 mg | TABLET EXEED-GM ✓ | Indications : Type 2 Diabetes. NIDDM With Vascular Complication. Dosage : Initiated Once Daily, and Gradually Titrated After Assessing Adequacy of Therapeutic Response. |
| 252. | Metformin 500 mg + Glibenclamide 5 mg + Pioglitazone 15 mg | TABLET EXEED-GMP ✓ | Indication : As 3rd line treatment of type II diabetes mellitus when diet, exercise and the single agents and the second line therapy with two drugs do not result in adequate glycemic control. Dosage: 1-2 tablet daily |
| 253. | Metformin 500 mg. + Gliclazide 80 mg. | TABLET EXEED-G ✓ | Indications : Patients with type 2 diabetes Dosage : One tablet daily. Max dose: 4 tablets |
| 254. | Metformin 500 mg. + Gliclazide 40 mg. + Pioglitazone 15 mg. | TABLET EXEED-GP | Indications : For the treatment of pulmonary arterial hypertension . Dosage : 5 mg once daily. May increase to 10 mg. if necessary. |
| 255. | Metformin 500 mg. + Gliclazide 80 mg. + Pioglitazone 15 mg. | TABLET EXEED-GP-PLUS ✓ | Indications : For the treatment of pulmonary arterial hypertension . Dosage : 5 mg once daily. May increase to 10 mg. if necessary. |
| 256. | Metformin 500 mg + Glipizide 5 mg | TABLET EXEED-Z-5 | Indications : Type 2 diabetes mellitus. Dosage : One tablet daily. Maximum recommended daily dose of 20 mg glipizide / 2000 mg metformin. |
| 257. | Metformin 500 mg. Glimepiride 1 mg. Atorvastatin 10 mg. | TABLET EXEED-AG-1 ✓ | Indications : For the treatment of type II diabetes mellitus associated with dyslipidemia. Dosage : 1 tablet daily which may be increased to 2 tablet daily. |
| 258. | Metformin 500 mg. Glimepiride 2 mg. Atorvastatin 10 mg. | TABLET EXEED-AG-2 ✓ | Indications : For the treatment of type II diabetes mellitus associated with dyslipidemia. Dosage : 1 tablet daily which may be increased to 2 tablet daily. |
| 259. | Metformin 500 mg + Pioglitazone 15 mg | TABLET EXEED-PG | Indications : Type 2 diabetes Dosage : One tablet daily. Max :The maximum recommended dose for pioglitazone is 30mg daily and for metformin is 2000 mg in adults |
| 260. | Metformin (ER) 1000 mg + Pioglitazone 15 mg | TABLET EXEED-PG-PLUS | Indication : Type 2 diabetes Mellitus Dosage : 1-2 tablets per day. |
| 261. | Metformin (ER) 1000 mg + Pioglitazone 30 mg | TABLET EXEED-PG-FORTE | Indication : Type 2 diabetes Mellitus Dosage : 1-2 tablets per day. |
| 262. | Nitrofurantoin 100 mg. (SR) | TABLET NITROFIT-100 ✓ | Indications : Treat infections of the bladder, Kidney & other parts of the Urinary Tract. Dosage : 50 - 100 mg. Max. dose : 4 times daily. |
| 263. | Pentosan Polysulfate Sodium 100 mg. | CAPSULE PPS-100 | Indications : For the relief of bladder pain or discomfort associated with interstitial cystitis Dosage : 100mg every 8 hours. Administration: 1 hour before or 2 hours after meals with water |
| 264. | Protein 60 gm. + Vitamins + Minerals + Spirulina + Ginseng + Mecobalamin & Other Essential Nutrients (Protein Powder Specially For Diabetic & Non-Diabetic & for All Age Group) | PROTEIN POWDER (200 GM) POWERME-SF SF=SUGAR FREE | Indications : Nutritional Supplement for diabetic & cardiac patients Dosage : Two tbls twice daily in fat free milk. |

ABBREVIATIONS :

DT = SWEET FLAVOURED DISPERSIBLE TABLET SR / CR / ER = SUSTAINED / CONTROLLED / EXTENDED RELEASE MD = SWEET FLAVOURED MOUTH DISSOLVING
 - AVAILABLE PRODUCTS ARE MARKED WITH TICK (✓) REST PRODUCTS ARE BEING MADE AVAILABLE ON MARKET DEMAND TIME TO TIME
 - BILAYERED = 2 COLOUR TABLET - (MEANS FRONT SIDE ONE COLOUR & BACK SIDE 2ND COLOUR)

| S.No. | FORMULA <small>{ Read Alphabetically }</small> | BRAND NAME | Where to use INDICATIONS and DOSAGE |
|----------------------|---|---|--|
| 265. | Pioglitazone 15 mg. | TABLET PGT -15 | Indication : Management of type 2 diabetes (NIDDM) Dosage : One tablet daily. Max -2 tablet |
| 266. 267. 268. | Repaglinide $\frac{0.5 \text{ mg.}}{1 \text{ mg.}}$ $\frac{1 \text{ mg.}}{2 \text{ mg.}}$ | TABLET REPAGOT $\frac{0.5}{1}$ $\frac{1}{2}$ | Indications : Type 2 (non-insulin dependent) diabetes. Dosage : Initially 0.5mg before each main meal. Maximum 4mg single dose before each main meal, maximum total daily dose 16mg. |
| 269. | Repaglinide 1 mg. + Metformin 500 mg. | TABLET REPAGOT-M1 | Indications : Type 2 diabetes Mellitus Dosage : 1 - 2 Tablets to taken with main meal. Max. 4 tab. per day |
| 270. | Repaglinide 2 mg. + Metformin 500 mg. | TABLET REPAGOT-M2 | Indications : Type 2 diabetes Mellitus Dosage : 1 - 2 Tablets to taken with main meal. Max. 4 tab. per day |
| 271. | Sevelamer Carbonate $\frac{400 \text{ mg.}}{800 \text{ mg.}}$ | TABLET SEVLOO $\frac{400}{800}$ | Indications : For the control of hyperphosphatemia in patients with end-stage renal disease (ESRD) undergoing dialysis Dosage : 1 tablet per day (400mg) to max 6 tablet per day (4.8 gm). Dose adjustments, when necessary should be done every 1 to 3 weeks by increasing one tablet per meal (3 tablets per day) until the desired serum levels are met. |
| 272. | Sildenafil Capsules 8 mg. | CAPSULE SIDOSIN-8 | Indications : treatment of the signs and symptoms of benign prostatic hyperplasia (BPH) Dosage : 1 tablets per day |
| 273. | Tamsulosin HCL 0.4 mg + Dutasteride 0.5 mg | TABLET TSD-9 ✓ | Indication : Benign prostatic Hyperplasia(BPH) Dosage : One tablet daily |
| 274. | Tamsulosin 0.4 mg.+ Finasteride 5 mg (MR) (MR = Modified Release) | TABLET TSF-9 ✓ | Indication : Benign Prostatic Hypertrophy. Dosage : 1 tablet once daily. |
| 275. | Tamsulosin HCl. (ER) 0.4 mg + Tolterodine Tartrate (ER) 4mg | TABLET TST-8 | Indications : For BPH with overactive bladder in adult only Dosage : 1 tablet daily |
| 276. | Teneligliptin 20 mg. | TABLET TINIGLIP-20 | Indications : Treatment of Type 2 Diabetes Mellitus. Dosage : 20 mg orally once daily. Max 40mg once daily |
| 277. | Teneligliptin + $\frac{20 \text{ mg.}}{500 \text{ mg.}}$ & $\frac{20 \text{ mg.}}{1000 \text{ mg.}}$ Metformin(ER) $\frac{500 \text{ mg.}}{1000 \text{ mg.}}$ {ER = Extended Release} | TABLET TINIGLIP-M $\frac{500 \text{ ER}}{1000 \text{ ER}}$ | Indications : Treatment of Type 2 Diabetes Mellitus. Dosage : 20 mg orally once daily. Max 40mg once daily (dose calculated on the bases of teneligliptin dosage.) |
| 278. | Trospium Chloride 20 mg. | TABLET TROFAST-20 | Indications : For the treatment of overactive bladder with the symptoms of urge urinary incontinence, urgency, and urinary frequency. Dosage : 1 tablet twice daily, max: 60mg daily. |
| 279. 280. | Ursodeoxycholic Acid $\frac{150 \text{ mg.}}{300 \text{ mg.}}$ | TABLET STONEFIX $\frac{150}{300}$ | Indications & Dosage : Gallstone Dissolution: 8-10 mg/kg/d In divided dose 8-12hr ; not to exceed 300 mg/dose . Maintenance: 250 mg at night X 6 months / Primary Biliary Cirrhosis:13-15 mg/kg/d in divided dose with food / Cystic Fibrosis Liver Disease |
| 281. | Vitamin D3 60000 IU | TABLET (CHEWABLE) GUDVIT-D3 <i>sugar free</i> ORANGE FLAVOURED | Indications : Osteomalacia, vitamin D deficiency, hyperparathyroidism, Osteoporosis, Vit. D dependent rickets. Dosage : Adult-min/max 200 - 60,000 I.U.; Pediatric :min/max 100 - 25,000 I.U. |

ABBREVIATIONS :

DT = SWEET FLAVOURED DISPERSIBLE TABLET SR / CR / ER = SUSTAINED / CONTROLLED / EXTENDED RELEASE MD = SWEET FLAVOURED MOUTH DISSOLVING
 - AVAILABLE PRODUCTS ARE MARKED WITH TICK (✓) REST PRODUCTS ARE BEING MADE AVAILABLE ON MARKET DEMAND TIME TO TIME
 - BILAYERED = 2 COLOUR TABLET - (MEANS FRONT SIDE ONE COLOUR & BACK SIDE 2ND COLOUR)

| S.No. | FORMULA { Read Alphabetically} | BRAND NAME | Where to use INDICATIONS and DOSAGE |
|--------------|--|--|---|
| 282. 283. | Voglibose 0.2 mg. 0.3 mg. | TABLET METBOSE 2/3 ✓ | Indication : To enhance glycaemic control Dosage : 0.2 - 0.3 mg three times daily. with each meal. |
| 284. | Voglibose 0.2 mg + Metformin 500 mg | TABLET METBOSE-M2 | Indication : Uncontrolled type 2 diabetes. Dosage : One tablet 3 times a day. the dosage however should be individualized. |
| 285. | Voglibose 0.3 mg + Metformin 500 mg (SR) | TABLET METBOSE-M3 ✓ BILAYER TABLETS | Indication : Uncontrolled type 2 diabetes. Dosage : One tablet 3 times a day. the dosage however should be individualized. |
| 286. | Aceclofenac 100 mg. + Paracetamol 325 mg. | TABLET MANGOFLAM ✓ | Indications : Ankylosing spondylitis; Osteoarthritis; Rheumatoid Arthritis, Dental pain, gynecological pain. Dosage : 1 tablet twice daily. |

Capsule

Sidosin-8
Silodosin Capsules 8 mg.

BILAYER TABLETS

TAKETEL-MTL **45/90**
Telmisartan 20 mg. & Metoprolol Succinate (ER) 25 mg. (TAKETEL-MTL-45)
Telmisartan 40 mg. & Metoprolol Succinate (ER) 50 mg. (TAKETEL-MTL-90)

TABLETS

GLIPTOP **50/100**
Vildagliptin 50 mg & 100 mg

TABLETS

GLIPTOP-M **500/1000**
Vildagliptin 50 mg and Metformin 500 mg
Vildagliptin 50 mg and Metformin 1000 mg

TABLETS

GLIPTOP-M **500 ER/1000 ER**
Vildagliptin 50 mg and Metformin 500 mg (Bilayered & Extended Release)
Vildagliptin 50 mg and Metformin 1000 mg (Bilayered & Extended Release)

TABLETS

DAPAGLIX-M **500 ER/1000 ER**
Dapagliflozin 20mg & Metformin 500mg (Bilayered & Extended Release)
Dapagliflozin 20mg & Metformin 1000mg (Bilayered & Extended Release)

TABLETS

TENEGLIP-M **500 ER/1000 ER**
Teneligliptine 20mg & Metformin 500mg (Bilayered & Extended Release)
Teneligliptine 20mg & Metformin 1000mg (Bilayered & Extended Release)

ABBREVIATIONS :

DT = SWEET FLAVOURED DISPERSIBLE TABLET SR / CR / ER = SUSTAINED / CONTROLLED / EXTENDED RELEASE MD = SWEET FLAVOURED MOUTH DISSOLVING
- AVAILABLE PRODUCTS ARE MARKED WITH TICK (✓) REST PRODUCTS ARE BEING MADE AVAILABLE ON MARKET DEMAND TIME TO TIME
- BILAYERED = 2 COLOUR TABLET - (MEANS FRONT SIDE ONE COLOUR & BACK SIDE 2ND COLOUR)

15

Perfect Remedies
DIABETIC & CARDIAC CARE